

pH IN PEST CONTROL: EFFECTS ON FORMULATION AND PERFORMANCE

Quadros S. F.¹ & Shetty S. S.²

¹SMark Corporation, Mumbai - 400054, India; ²Express Pesticides Pvt. Ltd., Mumbai- 400054, India

Email: stq@epcorn.com (Corresponding Author)

ABSTRACT

pH is a fundamental physicochemical parameter that influences nearly every stage of pest control product development and application, yet its role remains comparatively underrepresented in integrated discussions of formulation science and pest management. Beyond serving as a routine quality control measure, pH directly governs active ingredient stability, solubility, preservative efficiency, dispersion behaviour, emulsion formation, and storage performance, thereby affecting overall product efficacy. Variations in pH may accelerate degradation reactions, alter biological availability, and influence interactions between pest control products and environmental matrices. In formulation development, pH determines the compatibility of active compounds with surfactants, solvents, preservatives, and carrier systems. Improper pH adjustment may lead to precipitation, hydrolysis, phase separation, reduced shelf life, and loss of biological activity. During application, pH influences spray solution stability, water compatibility, residual performance, and persistence on treated surfaces. Environmental conditions including water hardness, soil chemistry, and surface characteristics further modify pH-dependent product behaviour. The importance of pH extends beyond formulation characteristics and may affect pest physiology through changes in feeding responses, microbial growth, sensory perception, and biochemical activity. Such interactions are increasingly relevant with the growth of botanical repellents, bio-based formulations, and environmentally responsive pest management technologies. This review consolidates current understanding of pH across formulation chemistry, application science, pest interactions, and environmental performance. The paper highlights existing knowledge gaps and discusses opportunities for improved formulation optimization and future development of more stable and effective pest control systems.

KEYWORDS: pH; Pest control; Formulation stability; Pesticide efficacy; Integrated pest management

INTRODUCTION

Effective pest control relies on the successful interaction of active ingredients, formulation components, application conditions, environmental factors, and biological responses of target organisms. While considerable emphasis has traditionally been placed on active ingredient

discovery and delivery technologies, comparatively less attention has been directed toward formulation variables that determine whether an active system performs as intended under practical conditions. Among these variables, pH represents one of the most influential yet frequently underestimated parameters affecting pest control outcomes.

pH, defined as the negative logarithm of hydrogen ion concentration, provides a measure of acidity or alkalinity within a system and serves as an indicator of chemical reactivity and biological compatibility. In agricultural, structural, urban, industrial, and public health pest management applications, pH influences multiple interconnected processes including active ingredient stability, solubility, emulsion behaviour, preservative performance, application efficiency, and environmental persistence.

The influence of pH on pest control products begins during formulation design. Commercial pest management products rarely consist solely of active ingredients; instead, they contain complex mixtures of solvents, surfactants, stabilizers, preservatives, emulsifiers, dispersants, fragrances, and carrier materials. Each component may contribute to the overall acid-base balance of the formulation and alter the behaviour of the final product [McClement & Jafari, 2018]. Consequently, pH optimization becomes necessary not only to preserve ingredient compatibility but also to maintain desired physicochemical characteristics throughout manufacturing and storage.

One of the most recognized consequences of inappropriate pH control is accelerated chemical degradation. Numerous pesticide classes exhibit pH-sensitive decomposition pathways, particularly hydrolysis and oxidation reactions. Organophosphate compounds, for example, frequently undergo rapid degradation under alkaline conditions, whereas pyrethroid compounds generally exhibit greater stability within slightly acidic environments. Such

degradation can substantially reduce active ingredient concentration before application or during field use, resulting in diminished pest control efficacy and increased treatment costs.

In addition to influencing chemical stability, pH affects physical formulation properties including solubility, suspension stability, droplet formation, and emulsion integrity. Minor shifts in pH may alter intermolecular interactions sufficiently to cause precipitation, flocculation, creaming, or phase separation [McClements, 2007]. These effects become increasingly significant in modern formulation systems such as suspension concentrates, microemulsions, nano-emulsions, encapsulated products, and botanical repellents where product performance depends heavily on controlled physicochemical behaviour.

The application stage introduces further complexity because product pH interacts with external variables including water quality, surface chemistry, temperature, and environmental exposure. Spray water characteristics, particularly alkalinity and hardness, may modify formulation pH after dilution and accelerate degradation processes. Surface pH can influence adsorption, residual activity, and persistence following application. Consequently, products optimized under laboratory conditions may perform differently under practical field environments.

Area	Main pH effect	Practical consequence
Active ingredients	Hydrolysis	Reduced efficacy
Emulsions	Instability	Phase separation
Preservatives	Reduced activity	Microbial growth
Spray solutions	Degradation	Field failure
Pest physiology	Biological response	Variable control

Table 1: Major areas influenced by pH in pest control

Recent interest in sustainable and low-toxicity pest management has further increased the relevance of pH optimization. Botanical formulations, microbial agents, and bio-based repellents frequently demonstrate greater sensitivity to pH variation than conventional synthetic chemistries. Essential oils and naturally derived compounds may undergo oxidation, hydrolysis, volatilization, or changes in sensory profile under unsuitable pH conditions, reducing product effectiveness and shelf stability.

Despite these broad implications, discussions of pH in pest control remain fragmented across formulation chemistry, application technology, environmental science, and pest biology literature.

Existing studies often evaluate pH as an isolated quality parameter rather than recognizing its integrated role across the entire product lifecycle.

Therefore, this review aims to provide a comprehensive overview of the role of pH in pest control by examining its effects on formulation stability, product performance, pest interactions, application conditions, and environmental behaviour. The review further identifies emerging opportunities for formulation optimization and highlights areas requiring additional investigation to improve the design of future pest management technologies.




Figure 1: Role of pH throughout the pest control lifecycle

FUNDAMENTALS OF pH IN PEST CONTROL

Concept and Measurement of pH

pH is a quantitative measure of hydrogen ion activity in a solution and is commonly used to describe the acidic or alkaline nature of chemical systems. Conventionally, pH is expressed as the negative logarithm of hydrogen ion concentration, where lower values indicate increased acidity and higher values indicate greater alkalinity. Although

pH is often treated as a simple numerical parameter, it functions as a broader indicator of chemical reactivity and strongly influences interactions among formulation ingredients and biological systems.

In pest control applications, pH serves as a critical determinant of product behaviour throughout formulation development, storage, application, and environmental exposure. Even relatively small changes in pH may substantially alter reaction

rates, molecular stability, and physical characteristics of a product. As a result, pH monitoring is routinely incorporated into formulation quality control, compatibility assessment, and stability testing procedures.

The practical significance of pH arises from its influence on ionization and equilibrium processes. Many formulation ingredients contain functional groups capable of gaining or losing hydrogen ions depending on surrounding conditions. These changes affect molecular charge, solubility, intermolecular attraction, and chemical susceptibility to degradation. Consequently, the same formulation may exhibit substantially different characteristics under different pH conditions [Akbari & Nour, 2018].

Measurement of pH in pest control systems is typically performed using potentiometric methods based on calibrated pH meters equipped with glass electrodes. Indicator strips and colorimetric methods may provide rapid field assessments but

generally lack the precision required for formulation development. Reliable pH determination requires consideration of temperature, sample preparation, ionic strength, and calibration procedures because these variables influence measured values.

For concentrated formulations and multiphase systems such as emulsions and suspensions, pH measurement may require standardized dilution methods to obtain representative values. Inadequate sampling or improper measurement conditions can lead to inaccurate conclusions regarding formulation performance and stability.

Sources of pH Variation in Pest Control Products

The pH of pest control products rarely originates from a single component and instead results from combined interactions among active ingredients and auxiliary formulation materials. Several variables contribute to final product pH and determine whether that value remains stable during storage and use.




Figure 2: Sources of pH variation in pest control formulations

Water quality represents one of the most significant external factors affecting pH. Water used during manufacturing or dilution may contain dissolved

minerals, bicarbonates, carbonates, and organic matter that shift the acid-base balance of the final system. Hard water containing elevated calcium

and magnesium concentrations frequently alters formulation behaviour and may indirectly accelerate instability reactions.

Active ingredients themselves contribute to pH variation depending on their chemical structure and dissociation behaviour. Certain compounds may produce acidic environments, while others demonstrate alkaline tendencies or undergo pH-dependent transformation reactions. Formulators therefore must evaluate ingredient compatibility across expected operating conditions.

Auxiliary ingredients including surfactants, emulsifiers, preservatives, solvents, and stabilizers further influence pH. Ionic surfactants may alter electrostatic interactions within formulations, whereas preservatives often display narrow functional pH ranges. Botanical extracts and essential oils may also contribute weakly acidic or oxidative components that influence formulation equilibrium over time.

Packaging and storage conditions can introduce additional variability. Exposure to temperature fluctuations, oxygen, light, and moisture may promote reactions that gradually shift formulation pH during shelf life. Consequently, pH is not merely a manufacturing specification but a dynamic property that requires monitoring throughout product storage and distribution.

Factor	Mechanism	Potential consequence
Water hardness	Alters alkalinity	Reduced stability
Surfactants	Changes ionic environment	Emulsion instability
Preservatives	Restricted pH activity	Microbial contamination
Temperature	Changes reaction rate	pH drift
Storage duration	Progressive degradation	Shelf-life reduction

Table 2: Typical factors affecting pH in pest control formulations

Relationship Between pH and Product Performance

The importance of pH in pest control extends beyond analytical measurement because it directly affects operational performance. Product stability, biological efficacy, physical appearance, and application consistency frequently depend on maintaining pH within an optimized range.

Chemical stability is among the most immediate consequences of pH variation. Hydrolysis reactions often accelerate under strongly acidic or alkaline conditions, reducing active ingredient concentration and shortening shelf life. Oxidation and polymerization reactions may also become more favourable outside optimal pH ranges.

Physical stability is similarly affected through changes in particle interaction and interfacial behaviour [Akbari & Nour, 2018]. Suspension systems may aggregate or settle, while emulsions may exhibit creaming, coalescence, or separation. Such changes influence dose uniformity and reduce application reliability.

Biological performance can also become pH dependent because formulation chemistry affects availability and interaction of active compounds with target organisms. Altered penetration, release behaviour, volatility, and persistence may modify the final pest control outcome despite unchanged nominal concentrations.

Taken together, these effects demonstrate that pH should not be considered an isolated laboratory parameter but rather an integrated design variable linking formulation chemistry to practical field performance.

ROLE OF PH IN PEST CONTROL FORMULATION SCIENCE

Formulation science is a critical component of modern pest control because the biological activity of an active ingredient depends not only on its intrinsic toxicity or repellency but also on the environment in which it is delivered. Commercial pest control products are highly engineered systems designed to optimize stability, usability, shelf life, and biological performance. Within these systems, pH acts as a central control parameter governing chemical compatibility, physical integrity, and application behaviour.

Unlike simple chemical mixtures, pest control formulations contain multiple interacting ingredients including active compounds, solvents, emulsifiers, surfactants, preservatives, stabilizers, rheology modifiers, fragrances, encapsulation materials, and carrier phases. The acid-base balance established among these components determines the final physicochemical characteristics of the product. Failure to maintain appropriate pH conditions may compromise performance long before the product reaches the target pest.

Consequently, pH optimization has become an essential component of formulation design and quality assurance in both conventional synthetic products and emerging bio-based pest management technologies.

Effect of pH on Active Ingredient Stability

Among all formulation variables, active ingredient stability is often the most directly influenced by pH. Chemical degradation reduces available active concentration and may generate by-products that alter efficacy, appearance, odour, or safety.

Hydrolysis is one of the most common degradation pathways affected by pH. Hydrolysis involves cleavage of chemical bonds through reaction with water and frequently accelerates under strongly acidic or alkaline conditions. Since many pest control products are water-based or become diluted before application, understanding hydrolytic behaviour is essential.

Several conventional pesticide classes exhibit marked pH sensitivity –

Organophosphate compounds are particularly susceptible to alkaline hydrolysis [Bokulić Petrić et. al., 2023; Zamy et. al., 2004]. Elevated pH may accelerate cleavage reactions and reduce active ingredient concentration during storage or after dilution. Such degradation may occur rapidly enough to reduce field efficacy if formulations remain exposed to unfavourable conditions for extended periods.

Carbamate compounds may also demonstrate pH-dependent decomposition, resulting in reduced residual activity and shorter effective treatment periods [Zamy et. al., 2004].

Pyrethroid compounds generally exhibit greater stability under mildly acidic conditions but may undergo increased degradation in alkaline environments [Camilleri, 1984; Takahashi et al., 1985]. Since pyrethroids are widely used in structural and public health pest control applications, pH control becomes especially relevant for maintaining residual effectiveness.

Botanical actives present additional challenges because many plant-derived compounds contain chemically reactive terpenes, aldehydes, esters, and phenolic constituents. These compounds may undergo oxidation, rearrangement, hydrolysis, or

aroma changes when exposed to unsuitable pH conditions.

Because degradation reactions follow kinetic behaviour, relatively small changes in pH may

produce disproportionately large reductions in shelf life and efficacy.




Figure 3: Effects of pH on active ingredient stability

Influence of pH on Solubility and Ingredient Availability

Solubility determines whether active ingredients remain uniformly distributed and available for biological action. pH directly affects molecular ionization and therefore alters dissolution behaviour and partitioning between formulation phases.

Many active compounds behave as weak acids or weak bases. Their ionization state changes depending on surrounding pH conditions, influencing water compatibility and phase distribution. Ionized molecules generally exhibit increased aqueous solubility but reduced affinity for hydrophobic environments, whereas non-ionized molecules may display opposite behaviour. Improper pH selection may produce incomplete dissolution, precipitation, or non-uniform distribution of active components. Such instability creates inconsistency in delivered dose and may reduce pest exposure.

This effect becomes particularly important in water-dispersible granules, suspension concentrates, soluble concentrates, and emulsifiable concentrates where ingredient distribution must remain stable over prolonged storage periods.

For botanical formulations, pH-dependent changes in oil-water interactions may influence release kinetics and sensory characteristics that contribute to repellent performance.

pH and Emulsion Stability

Emulsion-based systems represent one of the most widely used formulation approaches in pest control. These systems combine immiscible phases using surfactants that reduce interfacial tension and maintain dispersion.

The stability of emulsions is highly sensitive to pH because surface charge and interfacial interactions are controlled by ionization processes. Variations in

pH may alter surfactant behaviour and reduce electrostatic stabilization.

When emulsion stability decreases, several undesirable outcomes may occur including creaming, flocculation, coalescence, and complete phase separation [McClements, 2007]. Such instability reduces appearance quality, complicates dosing, and may compromise biological performance.

Acidic conditions may destabilize certain emulsifier systems through reduced charge

repulsion, while highly alkaline environments may promote hydrolysis of susceptible surfactants.

Modern pest control products increasingly employ advanced emulsion technologies including nano-emulsions and microemulsions [Marhamati et. al., 2021]. These systems provide improved dispersion and release control but frequently exhibit narrower acceptable pH ranges than conventional formulations [Safaya & Rotliwala, 2020].

Accordingly, optimization of pH is often integrated with surfactant selection and emulsion characterization during product development.




Figure 4: Influence of pH on emulsion and suspension systems

pH and Suspension Stability

Suspension formulations contain dispersed solid particles maintained within a liquid medium. Stability depends on controlling particle interactions and minimizing aggregation.

Changes in pH modify surface charge and zeta potential, influencing whether particles repel or attract one another. Insufficient electrostatic stabilization may produce flocculation and sedimentation.

Persistent sediment formation can create irreversible caking, making products difficult to

redisperse and reducing consistency during application.

Formulation attribute	pH influence	Practical outcome
Active stability	Hydrolysis	Reduced potency
Solubility	Ionization	Precipitation
Emulsion integrity	Surfactant performance	Separation
Suspension stability	Particle interactions	Sedimentation
Release behaviour	Diffusion kinetics	Variable efficacy

Table 3: Examples of formulation attributes influenced by pH

Control of pH therefore contributes not only to physical appearance but also to accurate delivery of

active ingredients and long-term formulation reliability.

Influence of pH on Preservative Performance and Microbial Stability

Preservation is an essential yet frequently underestimated aspect of pest control formulation science, particularly for water-based products and formulations containing natural ingredients. Microbial contamination during storage may reduce active ingredient concentration, alter product appearance, generate off-odours, and shorten commercial shelf life. Because preservative activity is strongly influenced by acid-base conditions, pH optimization becomes a necessary component of formulation stabilization.

Preservatives function through mechanisms that commonly involve membrane disruption, inhibition of enzyme systems, interference with nutrient transport, or suppression of microbial metabolism. However, many preservative compounds exhibit activity only within defined pH ranges because their antimicrobial effectiveness depends on molecular ionization.

Weak-acid preservatives frequently demonstrate improved performance under acidic conditions where larger proportions remain in non-ionized forms capable of penetrating microbial membranes [Russell, 2003]. As pH increases, ionization may reduce preservative penetration and lower antimicrobial activity.

This phenomenon becomes particularly relevant in botanical and low-solvent formulations because natural extracts often introduce nutrients or organic materials that increase microbial susceptibility. Water-rich formulations may therefore require simultaneous optimization of preservative

concentration and pH to maintain long-term stability.

Inadequate pH control may permit gradual microbial growth despite initially acceptable preservative levels. Such contamination may further modify formulation pH through metabolic activity, creating feedback mechanisms that accelerate product deterioration.

Consequently, preservative selection should be considered together with formulation pH rather than treated as an independent design decision.

pH and Shelf-Life Stability

Commercial pest control products are expected to maintain performance throughout storage, transportation, and use under variable environmental conditions. Shelf life therefore represents a key indicator of formulation quality.

The stability of a formulation over time reflects cumulative chemical and physical changes occurring under storage conditions. Since many degradation pathways exhibit pH dependence, pH serves as both a predictive and diagnostic parameter during stability evaluation.

Chemical instability may appear as active ingredient loss, discoloration, odour changes, precipitation, or viscosity shifts. Physical instability may include phase separation, sedimentation, crystallization, and changes in droplet or particle size.

Accelerated stability testing is commonly employed during product development to estimate long-term storage behaviour. Products are exposed to elevated temperatures and environmental stress conditions to identify likely failure mechanisms. Monitoring pH during these studies provides

insight into degradation trends and formulation robustness [Badrudodoza et. al., 2018].

A formulation exhibiting stable pH over time generally indicates greater resistance to internal reaction processes. Conversely, progressive pH drift may suggest ingredient incompatibility, preservative failure, oxidation reactions, or hydrolytic decomposition.

Accordingly, routine pH monitoring is widely integrated into manufacturing quality control and product release specifications.

Role of pH in Botanical and Bio-Based Pest Control Formulations

The growing demand for environmentally compatible pest control technologies has accelerated development of botanical repellents and biologically derived formulations. Compared with many conventional synthetic systems, these products often display increased sensitivity to pH variation.

Essential oils and plant-derived actives contain chemically diverse compounds including alcohols, aldehydes, ketones, esters, terpenes, and phenolics [Borges et. al., 2018]. These compounds may undergo oxidation, hydrolysis, rearrangement, polymerization, or volatilization depending on surrounding conditions.

Changes in pH may alter not only chemical stability but also odour profile and release characteristics, both of which influence repellency performance. Because many repellent products rely on controlled volatilization rather than direct toxicity, preservation of sensory characteristics becomes especially important.

Water-based botanical formulations present additional challenges because hydrophobic oils

must remain uniformly dispersed within aqueous environments. pH influences emulsifier behaviour, droplet stability, and interactions between oil and water phases.

Natural extracts may also contain pigments, organic acids, proteins, or trace minerals capable of changing formulation pH during storage. As a result, botanical formulations frequently require tighter pH control and more extensive compatibility testing than equivalent synthetic systems [Borges, et. al., 2018].

Future development of sustainable pest control products will likely increase emphasis on pH-responsive stabilization approaches.

pH in Encapsulation and Controlled-Release Technologies

Controlled-release technologies are increasingly used to improve persistence, reduce active ingredient loading, and enhance safety profiles. Encapsulation systems including microcapsules, polymer matrices, nano-emulsions, and liposomal carriers frequently exhibit pH-sensitive behaviour.

Encapsulation performance depends on maintaining structural integrity of carrier materials while allowing controlled release of active compounds. Because many encapsulating materials contain ionizable functional groups, environmental pH may influence swelling, permeability, diffusion rates, and degradation.

Controlled-release systems may be designed intentionally to respond to environmental pH changes, enabling selective activation under target conditions. Such approaches offer opportunities to reduce non-target exposure while maintaining treatment efficacy.

However, unintended pH shifts during manufacturing or storage may destabilize carrier systems and alter release profiles. Consequently, pH monitoring remains essential throughout product development and validation.

Manufacturing and Quality Control Considerations

From an industrial perspective, pH serves as both a formulation variable and a process control parameter. Manufacturing operations require reproducible conditions to ensure batch-to-batch consistency and regulatory compliance.

Process variables including ingredient addition sequence, mixing intensity, temperature, dilution procedures, and holding time may influence final product pH. Establishing acceptable operating windows helps reduce variability and improve production efficiency.

Quality control programs commonly include pH testing at multiple stages including raw material evaluation, in-process monitoring, finished product assessment, and shelf-life verification.

Rather than representing a final acceptance criterion alone, pH should be considered an integrated indicator of formulation health throughout the product lifecycle.

System	Primary concern
Emulsions	Phase separation
Suspensions	Sedimentation
Water-based repellents	Microbial growth
Botanical formulations	Oxidation and odour loss
Encapsulation systems	Release instability

Table 4: Formulation systems and typical pH-related concerns

pH AND PEST MANAGEMENT PERFORMANCE

While formulation quality establishes the potential effectiveness of a pest control product, actual field performance depends on how that formulation behaves under application and environmental conditions. Products that demonstrate acceptable laboratory stability may still perform inconsistently if pH changes occur during dilution, application, environmental exposure, or interaction with target organisms. Consequently, pH should be considered not only a formulation parameter but also an operational variable influencing pest management outcomes.

The transition from formulation to application introduces multiple sources of pH variability including spray water composition, environmental conditions, treated surface characteristics, and biological interactions. These variables determine whether the active system reaches and affects the target pest as intended.

Influence of pH on Spray Solutions and Water Quality

Water is one of the most important yet frequently overlooked components in pest control applications. Most liquid formulations undergo dilution before use, making the final spray solution substantially different from the original product.

Water chemistry varies according to geographical location and may contain dissolved bicarbonates, carbonates, calcium, magnesium, iron, and organic impurities [Bokulić Petrić et. al., 2023]. These constituents influence both pH and alkalinity, which in turn affect formulation behaviour after mixing.

High-pH spray solutions may accelerate degradation reactions, particularly alkaline hydrolysis [Fishel & Ferrell, 2007; Bokulić Petrić et. al., 2023]. Consequently, active ingredient concentration may decline during holding periods

before application. In operational settings where prepared solutions remain unused for extended periods, losses may become sufficiently large to influence treatment outcomes.




Figure 5: Relationship between spray water pH and pesticide degradation

Water hardness may further reduce effectiveness through interactions with surfactants and charged molecules. Such interactions can alter droplet behaviour, reduce dispersion efficiency, and increase formulation instability.

To minimize variability, many pest management programs recommend evaluating source water characteristics and using buffering or conditioning agents where necessary. Maintaining suitable pH after dilution can substantially improve consistency of field performance [Fishel & Ferrell, 2007; Bokulić Petrić et. al., 2023].

Because water quality differs among regions and facilities, application protocols should include pH assessment rather than assuming compatibility under all operating conditions.

Effect of pH on Surface Interaction and Residual Performance

Following application, pest control products interact with treated surfaces before contacting target organisms. These interactions strongly influence persistence, availability, and residual efficacy.

Surface properties including porosity, mineral composition, moisture content, and surface pH determine the extent of adsorption and chemical transformation. Porous or chemically active surfaces may alter local pH conditions and accelerate active ingredient degradation.

Concrete surfaces provide a commonly cited example because they often exhibit alkaline characteristics capable of influencing residual performance. Similar effects may occur on painted walls, masonry materials, metals, and treated wood.




Figure 6: Surface pH effects on residual performance

Surface-induced pH changes may alter molecular stability and reduce the amount of active ingredient remaining available for pest exposure. In repellent products, altered surface chemistry may also influence release rates and volatilization behaviour. Residual performance therefore depends not only on the formulation itself but also on maintaining compatibility between the product and the treated environment.

Understanding these interactions becomes increasingly important in structural pest management where products are expected to remain active for prolonged periods.

Influence of pH on Pest Physiology and Biological Response

Although pH is commonly discussed in relation to formulation chemistry, its influence extends into biological interactions with target organisms.

Many physiological processes in pests are regulated by local pH conditions. Digestive activity, enzyme function, sensory perception, membrane transport, and microbial associations may all respond to acid-base changes.

In insects, digestive environments often exhibit controlled pH ranges that optimize nutrient processing and enzymatic reactions. Alterations in formulation chemistry may influence active ingredient availability and uptake within these biological systems.

Contact formulations may also exhibit pH-dependent penetration through cuticular layers, thereby affecting delivered dose. Variations in ionization state can modify transport behaviour and alter biological effectiveness.

For repellent technologies, pH may influence release characteristics and sensory detection. Botanical formulations frequently depend on volatile compounds whose evaporation profiles and odour intensity are affected by formulation conditions.

Microbial pest management systems demonstrate additional sensitivity because microbial growth and viability are directly linked to environmental pH. Conditions that favour survival during storage may differ from those promoting field performance, creating formulation challenges.

These observations suggest that pH influences pest control outcomes through both direct biological effects and indirect changes in formulation behaviour.

pH and Environmental Persistence

Environmental persistence represents a critical determinant of both effectiveness and sustainability. Products that degrade too rapidly may fail to achieve adequate control, whereas excessive persistence may increase environmental exposure.

Environmental compartments including soil, water, and treated surfaces exhibit distinct pH conditions capable of modifying product behaviour after application.

Soil pH influences adsorption, mobility, degradation kinetics, and biological availability of applied compounds [Stangroom et. al., 2000]. Acidic and alkaline soils may therefore produce different persistence profiles for the same product.

Aquatic environments introduce additional complexity because pH influences hydrolysis, photochemical reactions, and interactions with dissolved constituents [Stangroom et. al., 2000]. Such effects become relevant in applications where runoff or environmental exposure may occur.

Environmental pH may also affect degradation products and transformation pathways, influencing both efficacy and ecological outcomes.

Accordingly, optimization of pest control systems increasingly requires consideration of environmental compatibility in addition to immediate biological performance.

pH as a Component of Integrated Pest Management

Integrated Pest Management (IPM) emphasizes coordinated use of chemical, biological, environmental, and operational approaches to achieve sustainable control. Within this framework, pH can be viewed as a supporting parameter that improves efficiency rather than as an isolated treatment factor.

Optimization of formulation pH may reduce active ingredient requirements by improving delivery efficiency and reducing degradation losses.

Water conditioning and compatibility assessment may improve treatment consistency.

Selection of products appropriate for local environmental conditions may extend residual performance.

Monitoring pH during preparation and application may contribute to more predictable pest control outcomes and reduce unnecessary retreatment.

As pest management continues moving toward lower chemical loads and increased precision, incorporation of pH-based optimization strategies may become increasingly important.

Therefore, pH management should be recognized as an enabling tool that supports formulation performance, environmental compatibility, and overall treatment success.

Stage	Influencing factor	Expected consequence
Dilution	Water alkalinity	Active degradation
Application	Surface chemistry	Reduced persistence
Exposure	Environmental pH	Variable efficacy
Biological contact	Uptake changes	Altered response

Table 5: Application-stage factors affecting pH performance

ENVIRONMENTAL, REGULATORY AND FUTURE PERSPECTIVES

Environmental Considerations of pH in Pest Control

Environmental sustainability has become a major consideration in modern pest management due to increasing concerns regarding ecological safety, non-target effects, and long-term environmental impacts. Within this context, pH influences not only product performance but also the

environmental behaviour of pest control agents after application.

The environmental fate of active ingredients is largely determined by degradation processes including hydrolysis, oxidation, photolysis, and microbial transformation. Since many of these reactions are pH dependent, environmental acidity or alkalinity can substantially alter persistence and mobility. Consequently, identical products may exhibit different environmental behaviours when applied under different soil or water conditions.




Figure 7: Environmental fate of pest control products under different pH conditions

Soil pH is particularly important because it affects adsorption to soil particles, movement through the soil profile, and availability for biological degradation. Acidic soils may increase adsorption of certain compounds, reducing mobility but potentially prolonging persistence. Conversely, alkaline soils may promote degradation or alter chemical speciation, affecting both efficacy and environmental exposure.

Aquatic systems present additional challenges because water pH directly influences chemical stability and transformation pathways. Changes in pH may affect degradation rates, toxicity profiles, and interactions with dissolved organic matter.

Such considerations are especially relevant in areas where pesticide runoff into aquatic ecosystems is possible.

In addition to synthetic pesticides, pH also affects the environmental behaviour of botanical products and biological control agents. Essential oils, microbial formulations, and naturally derived compounds may undergo pH-dependent degradation or transformation processes that influence persistence and effectiveness. Therefore, environmental pH should be considered during product development and environmental risk assessment.

Environmental compartment	Role of pH	Potential impact
Soil	Adsorption and degradation	Persistence changes
Surface water	Hydrolysis and transformation	Altered efficacy
Groundwater	Chemical mobility	Environmental exposure
Treated surfaces	Stability and residual activity	Performance variability

Table 6: Environmental effects of pH on pest control products

Regulatory Importance of pH Evaluation

Regulatory agencies worldwide recognize the importance of physicochemical properties in determining product quality, safety, and performance. Although pH alone does not define product effectiveness, it serves as an important indicator of formulation consistency and stability.

Product registration processes commonly require physicochemical characterization including pH determination, storage stability assessment, compatibility evaluation, and shelf-life studies. These data help demonstrate that a formulation maintains acceptable performance throughout its intended storage period.

Monitoring pH during product development can assist in identifying potential incompatibilities among formulation components before commercialization. Similarly, routine quality-control testing helps ensure that manufactured batches remain within established specifications.

International testing guidelines frequently include pH measurement as part of formulation evaluation protocols because pH changes may indicate chemical degradation, ingredient incompatibility, microbial contamination, or manufacturing variability [FAO, 2024]. Consequently, pH monitoring contributes to both product quality assurance and regulatory compliance.

As regulatory expectations increasingly emphasize product safety, sustainability, and performance consistency, the importance of comprehensive pH characterization is likely to expand further.

Emerging Technologies and Future Research Directions

The future of pest control formulation science is increasingly focused on improving efficacy while minimizing environmental impact. Within this evolving landscape, pH optimization offers numerous opportunities for innovation.

One promising area involves development of pH-responsive delivery systems capable of altering release behaviour under specific environmental conditions. Such systems may improve target specificity, reduce active ingredient requirements, and enhance persistence where needed.




Figure 8: Future pH-responsive pest control technologies

Advances in nanotechnology have created opportunities for pH-sensitive nano-emulsions, polymeric carriers, and encapsulated formulations [Safaya & Rotliwala, 2020]. These technologies may enable improved protection of active ingredients from degradation while providing controlled release following application.

Growing interest in botanical and biologically derived pest management products also highlights the need for additional research regarding pH stability and compatibility. Although botanical formulations are often promoted as environmentally friendly alternatives, relatively limited information is available regarding the influence of pH on their long-term storage performance, release characteristics, and field efficacy.

Future investigations should also focus on interactions among pH, microbial communities, and pest behaviour [Niu et. al., 2023]. Understanding how formulation chemistry influences biological responses may provide new opportunities for improving Integrated Pest Management programs.

Another important research area involves predictive modelling of pH-dependent stability [Niu et. al., 2023; Badruddoza et al., 2023]. Combining formulation science with computational approaches may help identify optimal formulation conditions more efficiently than traditional trial-and-error methods.

Finally, increased collaboration among formulation chemists, entomologists, microbiologists, toxicologists, and environmental scientists will be essential for developing a more comprehensive understanding of pH-related processes across the entire pest control lifecycle.

Research area	Current knowledge	Research need
Botanical repellents	Limited	Stability optimization
Nano-formulations	Emerging	Long-term performance
Microbial formulations	Moderate	Storage and field compatibility
Smart delivery systems	Early-stage	Commercial application
IPM integration	Limited	Practical implementation

Table 7: Research priorities for future pH studies in pest control

CONCLUSION

pH is one of the most influential physicochemical parameters affecting pest control products and their performance. Its effects extend far beyond simple measurement of acidity or alkalinity and encompass formulation stability, active ingredient preservation, emulsion behaviour, suspension integrity, preservative effectiveness, application performance, environmental persistence, and biological interactions.

Throughout the lifecycle of a pest control product, from formulation development to field application and environmental exposure, pH influences both chemical and physical processes that determine overall effectiveness. Improper pH conditions may accelerate degradation, reduce stability, alter release characteristics, and ultimately compromise pest management outcomes. Conversely, optimization of pH can improve product reliability, enhance efficacy, extend shelf life, and support sustainable pest management practices.

The growing adoption of botanical formulations, biological control products, nano-enabled delivery systems, and environmentally conscious pest management approaches further increases the importance of understanding pH-dependent behaviour. These emerging technologies often exhibit greater sensitivity to pH variation than conventional formulations and therefore require careful optimization during development.

Despite its recognized significance, pH remains an underappreciated factor in many areas of pest control research and practice. Future studies should adopt a more integrated perspective that considers pH as a connecting variable linking formulation

chemistry, application technology, pest biology, and environmental performance.

A greater emphasis on pH-based optimization has the potential to improve the effectiveness, consistency, and sustainability of next-generation pest management technologies.

REFERENCES

1. Akbari, S., & Nour, A. H. (2018). Emulsion types, stability mechanisms and rheology: A review. *International Journal of Innovative Research and Scientific Studies*, 1(1), 11–17. <https://doi.org/10.53894/ijirss.v1i1.4>
2. Badruddoza, A. Z. M., Yeoh, T., Shah, J. C., & Walsh, T. (2023). Assessing and predicting physical stability of emulsion-based topical semisolid products: A review. *Journal of Pharmaceutical Sciences*, 112(7), 1772–1793. <https://doi.org/10.1016/j.xphs.2023.03.014>
3. Bokulić Petrić, A., Stipičević, S., & Mešić, A. (2023). Stability of malathion, diazinon and chlorpyrifos in different water types: A review. *Journal of Central European Agriculture*, 24(4), 873–887. <https://doi.org/10.5513/JCEA01/24.4.3920>
4. Borges, D. F., Lopes, E. A., Moraes, A. R. F., Soares, M. S., Visôto, L. E., Oliveira, C. R., & Valente, V. M. M. (2018). Formulation of botanicals for the control of plant-pathogens: A review. *Crop Protection*, 110, 135–140. <https://doi.org/10.1016/j.cropro.2018.04.003>
5. Camilleri, P. (1984). Alkaline hydrolysis of some pyrethroid insecticides. *Journal of Agricultural and Food Chemistry*, 32(5), 1122–1124. <https://doi.org/10.1021/jf00125a048>

6. Fishel, F. M., & Ferrell, J. A. (2007). Water pH and the Effectiveness of Pesticides: PI-156/PI193, 6/2007. *EDIS*, 2007(17). <https://doi.org/10.32473/edis-pi193-2007>
7. Food and Agriculture Organization. (2024). *Hydrolysis rate: Pesticide registration toolkit*. Food and Agriculture Organization of the United Nations. <https://www.fao.org/pesticide-registration-toolkit/registration-tools/data-requirements-and-testing-guidelines/study-detail/en/c/1186906/>
8. Marhamati, M., Ranjbar, G., & Rezaie, M. (2021). Effects of emulsifiers on the physicochemical stability of Oil-in-water Nanoemulsions: A critical review. *Journal of Molecular Liquids*, 340, 117218. <https://doi.org/10.1016/j.molliq.2021.117218>
9. McClements D. J. (2007). Critical review of techniques and methodologies for characterization of emulsion stability. *Critical reviews in food science and nutrition*, 47(7), 611-649. <https://doi.org/10.1080/10408390701289292>
10. McClements, D. J., & Jafari, S. M. (2018). Improving emulsion formation, stability and performance using mixed emulsifiers: A review. *Advances in Colloid and Interface Science*, 251, 55-79. <https://doi.org/10.1016/j.cis.2017.12.001>
11. Niu, H., Wang, W., Dou, Z., Chen, X., Chen, X., Chen, H., & Fu, X. (2023). Multiscale combined techniques for evaluating emulsion stability: A critical review. *Advances in Colloid and Interface Science*, 311, 102813. <https://doi.org/10.1016/j.cis.2022.102813>
12. Russell A. D. (2003). Challenge testing: principles and practice. *International journal of cosmetic science*, 25(3), 147–153. <https://doi.org/10.1046/j.1467-2494.2003.00179.x>
13. Safaya, M., & Rotliwala, Y. C. (2020). Nanoemulsions: A review on low energy formulation methods, characterization, applications and optimization technique. *Materials Today: Proceedings*, 27, 454–459. <https://doi.org/10.1016/j.matpr.2019.11.267>
14. Stangroom, S. J., Collins, C. D., & Lester, J. N. (2000). Abiotic behaviour of organic micropollutants in soils and the aquatic environment. A review: II. Transformations. *Environmental Technology*, 21(8), 865–882. <https://doi.org/10.1080/09593332108618059>
15. Takahashi, N., Mikami, N., Matsuda, T., & Miyamoto, J. (1985). Hydrolysis of the pyrethroid insecticide cypermethrin in aqueous media. *Journal of Pesticide Science*, 10(4), 643–648. <https://doi.org/10.1584/jpestics.10.643>
16. Zamy, C., Mazellier, P., & Legube, B. (2004). Analytical and kinetic study of the aqueous hydrolysis of four organophosphorus and two carbamate pesticides. *International Journal of Environmental Analytical Chemistry*, 84(14–15), 1059–1068. <https://doi.org/10.1080/03067310412331298403>